



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

especial interest to readers of the MONTHLY because three of them are presidential addresses to the "Mathematical Association." The title of the book is "The organization of thought." It is published by Williams and Norgate, London.

The D. Van Nostrand Company has recently published a book on "Recreations in mathematics," by H. E. Licks. It has chapters on arithmetic, algebra, geometry, trigonometry, analytic geometry, calculus, astronomy and the calendar, mechanics and physics, and an appendix. It is not so extensive as the well-known "Mathematical recreations and essays" by W. W. R. Ball, but it contains considerable material not to be found in that standard work. The eight pages devoted to the cell of the honey bee will be new to many readers.

## PROBLEMS FOR SOLUTION.

SEND ALL COMMUNICATIONS ABOUT PROBLEMS TO B. F. FINKEL, Springfield, Mo.

**2689. Proposed by E. V. HUNTINGTON, Cambridge, Mass.**

Show that the maximum value of

$$y = \frac{\sin x \cos (x + \varphi)}{\cos (x + \theta) \sin (x + \varphi + \theta)}$$

is  $y_1 = (\cos \theta - \rho)/(\cos \theta + \rho)$ , where  $\rho = \sqrt{\sin^2 \varphi - \sin^2 \theta}$ .

This problem was suggested to the proposer by a professor of civil engineering, and has important applications in the theory of conjugate stresses.

*Note.*—It may facilitate the work to let  $\xi = 2x + \varphi + \theta$ .

**2690. Proposed by E. V. HUNTINGTON, Cambridge, Mass.**

Find the maximum value of

$$y = \frac{\sin x \cos (x + \varphi)}{\cos (x + \theta) \sin (x + \beta + \theta)}.$$

**2691. Proposed by ROGER A. JOHNSON, Hamline University.**

Show by purely geometric methods, without the use of the calculus, that the envelope of all circles whose centers are on a fixed circle and which touch a fixed diameter of that circle is a two-arched epicycloid. (Cf. Calculus problem, 423.)

**2692. Proposed by J. L. RILEY, Stephenville, Texas.**

A cube is cut at random by a plane, what is the chance that the section is a hexagon?

**2693. Proposed by W. F. HARLOW, Portland, Oregon.**

A cow is tethered with a rope, length  $l$ , to a peg on the opposite side of a wall, height  $h$ , the peg being at a distance  $a$  from the wall. Find the area over which the cow can graze.

**2694. Proposed by N. P. PANDYA, Sojitra, India.**

Find the locus of the centroid of a triangle, whose vertex lies on a given parabola, whose base of given length is a segment of a given straight line of unlimited length, and one of whose base angles is known.

**2695. Proposed by FRANK IRWIN, University of California.**

A positive number, which for convenience we will write as a fraction,  $a/b$ , is developed into a continued fraction,

$$a_1 - \frac{1}{a_2 - \frac{1}{a_3 - \dots}},$$

by the following process:

$$a = a_1 b - r_1, \quad b = a_2 r_1 - r_2, \quad r_1 = a_3 r_2 - r_3, \text{ etc.}$$

Here  $a_1, a_2, \dots$  are positive integers, and, if we suppose  $a, b$  to have been taken positive,  $b > r_1 > r_2 > \dots > 0$ . Show that if  $a/b$  be but slightly larger than a positive integer or zero, the numerators (as likewise the denominators) of the successive convergents will for a time be in arithmetical progression, and determine how long this phenomenon will continue.

**2696. Proposed by L. E. LUNN, Heron Lake, Minnesota.**

An air pipe 18 inches in diameter passes diagonally through a room from one lower corner to the opposite upper corner leaving through elliptical openings in the floor and ceiling, so that the ellipses are tangent to two boundaries of the floor and to the two opposite boundaries of the ceiling. If the room is  $10 \times 12 \times 8$ , find the remaining cubic capacity of the room.

**2697. Proposed by H. S. UHLER, Yale University.**

Show how to reduce the left-hand members of the following identities to their respective right members:

$$\sin^2(x + \tfrac{1}{2}y) - \sin(x + \tfrac{1}{2}y) \sin(x - \tfrac{1}{2}y) = \sin^2 y,$$

$$\sin(x + y) \sin(x + \tfrac{1}{2}y) - \sin x \sin(x + \tfrac{3}{2}y) = \sin \tfrac{1}{2}y \sin y,$$

$$\sin x \sin(x + \tfrac{1}{2}y) - \sin(x - \tfrac{1}{2}y) \sin(x + y) = \sin \tfrac{1}{2}y \sin y.$$

**2698. Proposed by WARREN WEAVER, Throop College of Technology, Pasadena, California.**

An urn contains  $N$  balls numbered from 1 to  $N$ . Of these  $n$  are drawn out and are arranged linearly according to the numbers on each. A certain ball is observed to be the  $k$ th in this line. What is the most probable number written on this ball?

## SOLUTIONS OF PROBLEMS.

**490 (Algebra). Proposed by HENRY HEATON, Atlantic, Iowa.**

Show that  $\sin 3^\circ = \frac{1}{16}(\sqrt{30} + \sqrt{10} - \sqrt{6} - \sqrt{2}) + \frac{1}{8}(\sqrt{5} + \sqrt{5} - \sqrt{15} + 3\sqrt{5})$ .

SOLUTION BY S. E. RASOR, The Ohio State University.

Since  $3^\circ = 12^\circ - 9^\circ$ ,  $12^\circ = 30^\circ - 18^\circ$ , and, for  $\theta = 18^\circ$ ,  $2\theta = 90^\circ - 3\theta$ , the sine and the cosine of  $9^\circ$ ,  $12^\circ$ ,  $18^\circ$ , and thus  $\sin 3^\circ$  may be found as follows:

We have

$$\begin{aligned} \sin 2\theta &= 2 \sin \theta \cos \theta = \sin(90^\circ - 3\theta) = \cos 3\theta = 4 \cos^3 \theta - 3 \cos \theta, \\ 4 \sin^2 \theta + 2 \sin \theta &= 1, \end{aligned}$$

and

$$\sin 18^\circ = \tfrac{1}{4}(\sqrt{5} - 1), \quad \cos 18^\circ = \tfrac{1}{4}\sqrt{10 + 2\sqrt{5}}.$$

Also from the identities,  $\sin \tfrac{1}{2}A \pm \cos \tfrac{1}{2}A = \pm \sqrt{1 \pm \sin A}$  for  $A = 18^\circ$ , we have

$$\sin 9^\circ = \tfrac{1}{4}(\sqrt{3} + \sqrt{5} - \sqrt{5} - \sqrt{5}), \quad \cos 9^\circ = \tfrac{1}{4}(\sqrt{3} + \sqrt{5} + \sqrt{5} - \sqrt{5}).$$

Also,

$$\sin 12^\circ = \sin(30^\circ - 18^\circ) = \sin 30^\circ \cos 18^\circ - \cos 30^\circ \sin 18^\circ = \tfrac{1}{8}(\sqrt{10 + 2\sqrt{5}} - \sqrt{15} + \sqrt{3}),$$

$$\cos 12^\circ = \tfrac{1}{8}(\sqrt{30} + 6\sqrt{5} + \sqrt{5} - 1).$$

Therefore,

$$\sin 3^\circ = \sin(12^\circ - 9^\circ) = \sin 12^\circ \cos 9^\circ - \cos 12^\circ \sin 9^\circ$$

$$\begin{aligned} &= \left( \frac{\sqrt{10 + 2\sqrt{5}} - \sqrt{15} + \sqrt{3}}{8} \right) \left( \frac{\sqrt{3} + \sqrt{5} + \sqrt{5} - \sqrt{5}}{4} \right) \\ &\quad - \left( \frac{\sqrt{30} + 6\sqrt{5} + \sqrt{5} - 1}{8} \right) \left( \frac{\sqrt{3} + \sqrt{5} - \sqrt{5} - \sqrt{5}}{4} \right) \end{aligned}$$